

BINARY ADDITIVE PROBLEMS: RECURSIONS FOR NUMBERS OF REPRESENTATIONS

VLADIMIR SHEVELEV

ABSTRACT. We prove some general recursions for the numbers of representations of positive integers as a sum $x + y, x \in X, y \in Y$, where X, Y are increasing sequences. In particular, we obtain recursions for the number of the Goldbach, Lemoine-Levy, Chen and other binary partitions.

1. INTRODUCTION AND MAIN RESULTS

Let $S = \{s_n\}_{n \geq 1}, T = \{t_n\}_{n \geq 1}$ be increasing sequences (finite or infinite) of positive odd numbers and $W = \{w_n\}_{n \geq 1}$ be increasing sequence of the common terms of sequences S and T . In this case we are writing $W = S \cap T$. Denote $S(x), T(x)$ and $W(x)$ the corresponding counting functions of these sequences, e.g., $W(x)$ denotes $\#\{n : w_n \leq x\}$. For a positive even integer x , denote $g(x)$ the number of solutions of the diophantine equation

$$(1) \quad x = s + t, \quad s \in S, t \in T,$$

when the order of the summands is not important; in particular, in the case of $s, t \in W$ we have only solution corresponding to s, t . Everywhere below $s \in S, t \in T, w \in W$, etc. In Section 2 we prove the following recursion.

Theorem 1. *(Number of decompositions of even numbers into two odd numbers).*

For even $x \geq 4$,

$$g(x) = \sum_{t_1 \leq t \leq x/2} S(x-t) + \sum_{s_1 \leq s \leq x/2} T(x-s) - \sum_{w_1 \leq w \leq x/2} W(x-w)$$

$$(2) \quad - S(x/2)T(x/2) + \binom{W(x/2)+1}{2} - g(x-2) - g(x-4) - \dots - g(2).$$

Corollary 1. *If $S \subseteq T$, then, for even $x \geq 4$,*

$$g(x) = \sum_{t_1 \leq t \leq x/2} S(x-t) + \sum_{s_1 \leq s \leq x/2} (T(x-s) - S(x-s))$$

$$-S(x/2)T(x/2) + \binom{S(x/2)+1}{2} - g(x-2) - g(x-4) - \dots - g(2).$$

Corollary 2. *If $S = T$, then, for even $x \geq 4$,*

$$g(x) = \sum_{s_1 \leq s \leq x/2} S(x-s) - \binom{S(x/2)}{2} - g(x-2) - g(x-4) - \dots - g(2).$$

Let, furthermore, $L = \{l_n\}_{n \geq 1}, M = \{m_n\}_{n \geq 1}$ be increasing sequences of positive even numbers and $Z = \{z_n\}_{n \geq 1} = L \cap M$ and $L(x), M(x)$ and $Z(x)$ are their corresponding counting functions. For a positive even integer x , denote $e(x)$ the number of solutions of the diophantine equation

$$x = l + m, \quad l \in L, m \in M,$$

when the order of the summands is not important.

Theorem 2. *(Number of decompositions of even numbers into two even numbers).*

For even $x \geq 2$,

$$e(x) = \sum_{m_1 \leq m \leq x/2} L(x-m) + \sum_{l_1 \leq l \leq x/2} M(x-l) - \sum_{z_1 \leq z \leq x/2} Z(x-z) \\ - L(x/2)M(x/2) + \binom{Z(x/2)+1}{2} - e(x-2) - e(x-4) - \dots - e(0).$$

Corollary 3. *If $L \subseteq M$, then, for even $x \geq 2$,*

$$e(x) = \sum_{m_1 \leq m \leq x/2} L(x-t) + \sum_{l_1 \leq l \leq x/2} (M(x-s) - M(x-s)) \\ - L(x/2)M(x/2) + \binom{L(x/2)+1}{2} - e(x-2) - e(x-4) - \dots - e(0).$$

Corollary 4. *If $L = M$, then, for even $x \geq 2$,*

$$e(x) = \sum_{l_1 \leq l \leq x/2} L(x-l) - \binom{L(x/2)}{2} - e(x-2) - e(x-4) - \dots - e(0).$$

Finally, let $U = \{u_n\}_{n \geq 1}$ ($V = \{v_n\}_{n \geq 1}$) be increasing sequence of positive even (odd) numbers, and $U(x)$ ($V(x)$) be its counting function.

Theorem 3. *(Number of decompositions of odd numbers into one even and one odd numbers).*

Let $h(x)$, for a positive odd integer x , denote the number of solutions of the diophantine equation

$$(3) \quad x = u + v, \quad u \in U, v \in V.$$

Then, for $x \geq 3$,

$$(4) \quad \begin{aligned} h(x) = & \sum_{v_1 \leq v \leq (x+1)/2} U(x-v) + \sum_{u_1 \leq u \leq (x+1)/2} V(x-u) \\ & - U((x+1)/2)V((x+1)/2) - h(x-2) - h(x-4) - \dots - h(1). \end{aligned}$$

In Section 2 we prove Theorems 1 - 3; in Section 3 we give some examples including recursions for numbers of the Goldbach, Lemoine-Levy, Chen and other binary partitions.

2. PROOF OF THEOREMS 1 - 3

Proof of Theorem 1. a) We start with the base of induction of formula (2). Put in (2) $x = 4$. Let us prove that (2) is valid for every values of parameters s_1, t_1, w_1 . For $x = 4$, denoting right hand side of (2) via r , we have

$$(5) \quad \begin{aligned} r = & \sum_{t_1 \leq t \leq 2} S(4-t) + \sum_{s_1 \leq s \leq 2} T(4-s) - \sum_{w_1 \leq w \leq 2} W(4-w) \\ & - S(2)T(2) + \binom{W(2)+1}{2} - g(2). \end{aligned}$$

We distinguish several cases:

1) $s_1 = 1, t_1 = 1$. Then also $w_1 = 1$. Evidently, $g(2) = 1$. By (5), we have

$$r = S(3) + T(3) - W(3) - 1.$$

1a) $s_2 = 3, t_2 = 3$, then $w_2 = 3$. Therefore, $r = 1$. Evidently, we have only representation $4=3+1$, and thus $r = g(4)$.

1b) $s_2 = 3, t_2 \geq 5$, then $w_2 \geq 5$. Again $r = 1$ and $r = g(4)$.

1c) Symmetrically in case of $t_2 = 3, s_2 \geq 5$ we have $r = g(4) = 1$.

1d) $s_2 \geq 5, t_2 \geq 5$, then $w_2 \geq 5$ and $r = 0$. Evidently, $g(4) = 0$ and again $r = g(4)$.

2) $s_1 = 1, t_1 = 3$, then $w_1 \geq 3$ and $g(2) = 0$. Now, according to (5), we have

$$r = T(3) = 1 = g(4).$$

3) Symmetrically in case of $t_1 = 1, s_1 = 3$ we have $r = S(3) = 1 = g(4)$.

4) Finally, if $s_1 \geq 3, t_1 \geq 3$ then $w \geq 3$ and $g(2) = 0$. Now, by (5), $r = 0$ which corresponds to $g(4) = 0$.

Thus the base of induction is valid.

b)Step of induction. Denote

$$\begin{aligned}
 G(x) &= \sum_{t_1 \leq t \leq x/2} S(x-t) + \sum_{s_1 \leq s \leq x/2} T(x-s) - \sum_{w_1 \leq w \leq x/2} W(x-w) \\
 (6) \quad &- S(x/2)T(x/2) + \binom{W(x/2) + 1}{2}.
 \end{aligned}$$

We should prove that

$$(7) \quad G(x+2) - G(x) = g(x+2).$$

We have

$$\begin{aligned}
 G(x+2) - G(x) &= \sum_{t_1 \leq t \leq x/2+1} (S(x+2-t) - S(x-t)) \\
 &+ \sum_{s_1 \leq s \leq x/2+1} (T(x+2-s) - T(x-s)) - \sum_{w_1 \leq w \leq x/2+1} (W(x+2-w) - W(x-w)) \\
 &+ \begin{cases} 0, & \text{if } x/2+1 \notin S \cup T, \\ T(x/2-1), & \text{if } x/2+1 \in S \setminus W, \\ S(x/2-1), & \text{if } x/2+1 \in T \setminus W, \\ S(x/2-1) + T(x/2-1) - W(x/2-1), & \text{if } x/2+1 \in W \end{cases} \\
 (8) \quad &-S(x/2+1)T(x/2+1) + S(x/2)T(x/2) + \binom{W(x/2+1) + 1}{2} - \binom{W(x/2) + 1}{2}.
 \end{aligned}$$

First of all, notice that if $x/2+1 \in S \cup T$ then $x/2$ is even and, consequently, $x/2 \notin S \cup T$. Therefore, we can rewrite (8) as follow:

$$\begin{aligned}
 G(x+2) - G(x) &= \sum_{t_1 \leq t \leq x/2+1} (S(x+2-t) - S(x-t)) \\
 &+ \sum_{s_1 \leq s \leq x/2+1} (T(x+2-s) - T(x-s)) - \sum_{w_1 \leq w \leq x/2+1} (W(x+2-w) - W(x-w)) \\
 &+ \begin{cases} 0, & \text{if } x/2+1 \notin S \cup T, \\ T(x/2), & \text{if } x/2+1 \in S \setminus W, \\ S(x/2), & \text{if } x/2+1 \in T \setminus W, \\ S(x/2) + T(x/2) - W(x/2), & \text{if } x/2+1 \in W \end{cases} \\
 (9) \quad &-S(x/2+1)T(x/2+1) + S(x/2)T(x/2) + \binom{W(x/2+1) + 1}{2} - \binom{W(x/2) + 1}{2}.
 \end{aligned}$$

Now we verify directly that

$$S(x/2+1)T(x/2+1) - S(x/2)T(x/2) =$$

$$(10) \quad \begin{cases} 0, & \text{if } x/2 + 1 \notin S \cup T, \\ T(x/2), & \text{if } x/2 + 1 \in S \setminus W, \\ S(x/2), & \text{if } x/2 + 1 \in T \setminus W, \\ S(x/2) + T(x/2) + 1, & \text{if } x/2 + 1 \in W \end{cases}$$

and that

$$(11) \quad \binom{W(x/2 + 1) + 1}{2} - \binom{W(x/2) + 1}{2} = W(x/2) + 1.$$

From (9)-(11) we find

$$\begin{aligned} G(x+2) - G(x) &= \sum_{t_1 \leq t \leq x/2+1} (S(x+2-t) - S(x-t)) \\ &+ \sum_{s_1 \leq s \leq x/2+1} (T(x+2-s) - T(x-s)) - \sum_{w_1 \leq w \leq x/2+1} (W(x+2-w) - W(x-w)) = \\ (12) \quad &\sum_{t_1 \leq t \leq x/2+1: x+2-t \in S} 1 + \sum_{s_1 \leq s \leq x/2+1: x+2-s \in T} 1 - \sum_{w_1 \leq w \leq x/2+1: x+2-w \in W} 1. \end{aligned}$$

Notice that, if $s \leq x/2 + 1$, then $t = x + 2 - s \geq x/2 + 1$, if $t \leq x/2 + 1$, then $s = x + 2 - t \geq x/2 + 1$, and if $s = x/2 + 1$, then $t = x + 2 - s = x/2 + 1 \in W$, $t = x/2 + 1$, then $s = x + 2 - t = x/2 + 1 \in W$. Therefore, sum (12), indeed, is $g(x+2)$. ■

Proof of Theorem 2 is quite analogous to proof of Theorem 1.

Proof of Theorem 3. a) Put in (4) $x = 3$. Let us prove that (4) is valid for every values of parameters u_1, v_1 . For $x = 3$, denoting right hand side of (4) via r , we have

$$(13) \quad r = \sum_{v_1 \leq v \leq 2} U(3-v) + \sum_{u_1 \leq u \leq 2} V(3-u) - U(2)V(2) - h(1).$$

We distinguish several cases:

1) $u_1 = 0, v_1 = 1$. Then $h(1) = 1, V(2) = 1$ and, by (13), we have

$$(14) \quad r = V(3) - 1 + \begin{cases} 1, & \text{if } u_2 = 2, \\ 0 & \text{if } u_2 \geq 4. \end{cases}$$

1a) $u_2 = 2, v_2 = 3$, then $V(3) = 2$ and by (14) we have $r = 2$. In this case we have two representations of 3: $3=0+3$ and $3=2+1$. Thus $r = h(3)$.

1b) $u_2 = 2, v_2 \geq 5$, then $V(3) = 1$ and, by (14), $r = 1$. Here there is only representation: $3=2+1$ and again $r = h(3)$.

1c) $u_2 \geq 4, v_2 = 3$, then $V(3) = 2$ and, by (14), $r = 1$. Here there is only representation: $3=0+3$ and again $r = h(3)$.

1d) $u_2 \geq 4, v_2 \geq 5$, then $V(3) = 1$ and ,by (14), $r = 0$. Evidently, in this case $h(3) = 0$ as well.

2) $u_1 = 2, v_1 = 1$. Then $h(1) = 0, V(2) = U(2) = 1$ and, by (13), we have $r = V(1) = 1$. Evidently, in this case $h(3) = 1$ as well.

3) $u_1 = 0, v_1 = 3$. Then $h(1) = 0, V(2) = U(2) = 1$ and, by (13), we have $r = V(1) = 1$. Evidently, in this case $h(3) = 1$ as well.

4) $u_1 \geq 4, v_1 \geq 1$ or $u_1 \geq 0, v_1 \geq 5$, then evidently $r = h(3) = 0$. b)Step of induction. Denote

$$\begin{aligned} H(x) = & \sum_{v_1 \leq v \leq (x+1)/2} U(x-v) + \sum_{u_1 \leq u \leq (x+1)/2} V(x-u) \\ (15) \quad & - U((x+1)/2)V((x+1)/2). \end{aligned}$$

We should prove that

$$(16) \quad H(x+2) - H(x) = h(x+2).$$

We have

$$\begin{aligned} H(x+2) - H(x) = & \sum_{v_1 \leq v \leq (x+1)/2} (U(x+2-v) - U(x-v)) \\ & + \sum_{u_1 \leq u \leq (x+1)/2} (V(x+2-u) - V(x-u)) \\ & + \begin{cases} 0, & \text{if } (x+3)/2 \notin U \cup V, \\ U((x+1)/2), & \text{if } (x+3)/2 \in V, \\ V(x+1)/2, & \text{if } (x+3)/2 \in U \end{cases} \\ (17) \quad & - U((x+3)/2)V((x+3)/2) + U((x+1)/2)V((x+1)/2). \end{aligned}$$

Now, taking into account that $U \cap V = \emptyset$, we verify directly that

$$\begin{aligned} & U((x+3)/2)V((x+3)/2) - U((x+1)/2)V((x+1)/2) = \\ & + \begin{cases} 0, & \text{if } (x+3)/2 \notin U \cup V, \\ U((x+1)/2), & \text{if } (x+3)/2 \in V, \\ V(x+1)/2, & \text{if } (x+3)/2 \in U \end{cases} \end{aligned}$$

and from (17) we find

$$\begin{aligned} H(x+2) - H(x) = & \sum_{v_1 \leq v \leq (x+1)/2} (U(x+2-v) - U(x-v)) \\ & + \sum_{u_1 \leq u \leq (x+1)/2} (V(x+2-u) - V(x-u)) = \end{aligned}$$

$$(18) \quad \sum_{v_1 \leq v \leq (x+1)/2: x+2-v \in U} 1 + \sum_{u_1 \leq u \leq (x+1)/2: x+2-u \in V} 1.$$

Notice that, if $v \leq (x+1)/2$, then $u = x+2-v > (x+1)/2$; if $u \leq (x+1)/2$, then $v = x+2-u > (x+1)/2$. Therefore, sum (18) is, indeed, $h(x+2)$. ■

3. SOME EXAMPLES

Example 1. *Number of decompositions of $2n$ ($n \geq 1$) into unordered sums of two odd primes (Goldbach partitions)*

The first terms of this sequence are (see A002375 in [1]):

$$(19) \quad 0, 0, 1, 1, 2, 1, 2, 2, 2, 2, 3, 3, 3, 2, 3, 2, 4, 4, 2, 3, 4, 3, 4, 5, 4, 3, 5, 3, 4, 6, \dots$$

Let $\pi(x)$ be the counting prime function, $\pi_1(x)$ be the counting function of odd primes. Note that, Hardy and Wright give a simple explicit formula for $\pi(n)$ for $n \geq 4$ (see [2]):

$$\pi(n) = -1 + \sum_{j=3}^n ((j-2)! - j \lfloor (j-2)!/j \rfloor).$$

Denote $P = \{p\}$ the set of all primes, P_1 the set of odd primes. Putting in Corollary 2 $x = 2n$ and $S = P_1$, we have a recursion

$$(20) \quad g(2n) = \sum_{3 \leq p \leq n} \pi_1(2n-p) - \binom{\pi_1(n)}{2} - g(2n-2) - g(2n-4) - \dots - g(2), \quad n \geq 2, \quad g(2) = 0.$$

Example 2. *Number of decompositions of $2n$ ($n \geq 1$) into sums of a prime and a prime or semiprime (Chen partitions)*

Let $P_2 = \{p_1 p_2\}$ be the set of semiprimes, i.e. products of two (possibly equal) primes. Denote $P_{2,1} = \{p_{2,1}\}$ the subset of odd semiprimes. If $\pi_2(x)$ is the counting function of P_2 , then, by Noel-Panos-Wilson formula (see [2]), we have

$$(21) \quad \pi_2(x) = \sum_{p \leq \sqrt{x}, p \in P} (\pi(x/p) - \pi(p) + 1).$$

Denoting $\pi_{2,1}(x)$ the counting function of set $P_{2,1}$, by (21) we have

$$(22) \quad \pi_{2,1}(x) = \sum_{3 \leq p \leq \sqrt{x}, p \in P} (\pi(x/p) - \pi(p) + 1).$$

Putting in Corollary 1 $S = P_1, T = P_1 \cup P_{2,1}, x = 2n$, we obtain a recursion for the numbers of Chen "odd-odd" partitions:

$$\begin{aligned}
 g_1(2n) = & \sum_{9 \leq p_{2,1} \leq n} \pi_1(2n - p_{2,1}) + \sum_{3 \leq p \leq n} \pi_{2,1}(2n - p) - \pi_1(n)(\pi_{2,1}(n) + \pi_1(n)/2 - 1/2) \\
 (23) \quad & - g_1(2n - 2) - \dots - g_1(2), \quad n \geq 2, g_1(2) = 0.
 \end{aligned}$$

The first terms of this sequence are:

$$(24) \quad 0, 0, 1, 1, 2, 2, 3, 3, 3, 4, 5, 4, 6, 6, 4, 6, 6, 6, 8, 7, 7, \dots$$

If to take into account that the number of Chen "even-even" partitions is evidently

$$g_2(2n) = \begin{cases} 1, & \text{if } n - 1 \in P \cup \{1\}, \\ 0, & \text{if } n - 1 \notin P \cup \{1\} \end{cases},$$

then the first terms of all Chen partitions are:

$$(25) \quad 0, 1, 2, 2, 2, 3, 3, 4, 3, 4, 5, 5, 6, 7, 4, 6, 6, 7, 8, 8, 7, \dots$$

Example 3. *Number of decompositions of $2n - 1$ ($n \geq 1$) into sums of a prime and a doubled prime (Lemoine-Levy partitions)*

Put in Theorem 3 $U = 2P, V = P, x = 2n - 1, n \geq 2$. Then $U(x) = \pi(x/2), V(x) = \pi(x)$ and we find a recursion for required numbers:

$$\begin{aligned}
 h(2n - 1) = & \sum_{2 \leq p \leq n} \pi(n - (p + 1)/2) + \sum_{2 \leq p \leq \lfloor n/2 \rfloor} \pi(2(n - p) - 1) - \pi(n)\pi(n/2) \\
 (26) \quad & - h(2n - 3) - \dots - h(1), \quad n \geq 2, h(1) = 0.
 \end{aligned}$$

The first terms of this sequence are (see A046927 in [1]):

$$(27) \quad 0, 0, 0, 1, 2, 2, 2, 2, 4, 2, 3, 3, 3, 4, 4, 2, 5, 3, 4, 4, 5, 4, 6, 4, 4, 7, \dots$$

Example 4. *Number of decompositions of $4n + 1$ ($n \geq 0$) into unordered sums of two squares of nonnegative integers and number of decompositions of n ($n \geq 0$) into unordered sums of two triangular numbers*

Let $Q_1 = \{q_1\} (Q_2 = \{q_2\})$ be the set of odd (even) squares of integers. Then its counting function is $\lfloor (\sqrt{x} + 1)/2 \rfloor$ ($\lfloor (\sqrt{x} + 2)/2 \rfloor$). Putting in Theorem 3 $U = Q_2, V = Q_1, x = 4n + 1$ and taking into account that $h(4i - 1) = 0, i \geq 1$, we obtain the following recursion:

$$\begin{aligned}
 h(4n + 1) = & \sum_{0 \leq q_2 \leq 2n} \lfloor (1 + \sqrt{4n - q_2 + 1})/2 \rfloor + \sum_{1 \leq q_1 \leq 2n + 1} \lfloor (2 + \sqrt{4n - q_1 + 1})/2 \rfloor \\
 & - \lfloor (1 + \sqrt{2n + 1})/2 \rfloor \lfloor (2 + \sqrt{2n + 1})/2 \rfloor - h(4n - 3) - h(4n - 7) - \dots - h(1), \\
 (28) \quad & n \geq 1, h(1) = 1.
 \end{aligned}$$

With help of the following statement we shall give a simpler recursion.

Lemma 1. *For $n \geq 0$, the number $4n + 1$ is a sum of two squares if and only if n is a sum of two triangular numbers. Moreover, the number of unordered representations of $4n + 1$ as a sum of two squares of nonnegative integers equals to the number of unordered representations of n as a sum of two triangular numbers.*

Proof. For integer $x \geq 0$, denote $T_x = x(x + 1)/2$. If $n = T_x + T_y$, then one can verify directly that

$$4n + 1 = (x + y + 1)^2 + (x - y)^2.$$

Conversely, if $4n + 1 = x^2 + y^2$, then one can verify that

$$n = T_{(x+y-1)/2} + T_{(|x-y|-1)/2}. \blacksquare$$

Remark 1. *Notice that, the identity*

$$4T_x + 4T_y + 1 = (x + y + 1)^2 + (x - y)^2$$

is a generalization of well known identity $T_x + T_{x-1} = x^2$.

Let now $L = \{l\}_{l \geq 0}$, where $l = j(j + 1)$, $j \geq 0$. Note that, $L(x) = \lfloor (1 + \sqrt{4x + 1})/2 \rfloor$. Consider representations of the form $2n = l_1 + l_2$, $l_1, l_2 \in L$, $n \geq 0$ or, equivalently, $n = T_x + T_y$. The number $t(n) = e(2n)$ of such representations we find by Corollary 4 for $x = 2n$. After small transformation we obtain the following recursion:

$$(29) \quad t(n) = \sum_{k=1}^{\lfloor (1 + \sqrt{4n+1})/2 \rfloor} \left\lfloor (1 + \sqrt{1 + 4(2n - k^2 + k)})/2 \right\rfloor - \binom{\lfloor (1 + \sqrt{4n+1})/2 \rfloor}{2} - t(n-1) - t(n-2) - \dots - t(0), \quad n \geq 1, \quad t(0) = 1.$$

By Lemma 1,

$$h(4n + 1) = t(n).$$

The first terms of this sequence are (see A052343 in [1]):

$$(30) \quad 1, 1, 1, 1, 1, 0, 2, 1, 0, 1, 1, 1, 1, 0, 1, 2, 0, 1, 0, 1, 2, 1, 0, 1, 1, \dots$$

REFERENCES

- [1] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences* (<http://www.research.att.com>)
- [2] E. W. Weisstein, "Goldbach Partition", "Prime Counting Function", "Semiprime" From *MathWorld—A Wolfram Web Resource* (<http://mathworld.wolfram.com/GoldbachPartition.html>)